

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 023 834 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
02.08.2000 Bulletin 2000/31

(21) Application number: 00300637.6

(22) Date of filing: 28.01.2000

(51) Int. Cl.⁷: **A01N 35/04**

// (A01N35/04, 59:20, 59:02,
55:00, 47:44, 47:38, 47:14,
47:04, 43:90, 43:84, 43:82,
43:76, 43:653, 43:60, 43:56,
43:54, 43:42, 43:40, 43:36,
43:32, 43:30, 37:50, 37:46,
37:38, 37:34, 37:06)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 29.01.1999 US 240412

(71) Applicant:
**American Cyanamid Company
Madison, New Jersey 07940-0874 (US)**

(72) Inventors:
• **Cotter, Henry Van Tuyl
Trenton, NJ 08618 (US)**


• **Reichert, Gunter
55270 Bubenheim (DE)**
• **Sieverding, Ewald
55578 St. Johann (DE)**
• **Jegerings, Petrus Martinus Franciscus Emanuel
1300 Wavre (BE)**

(74) Representative:
**Walters, Philip Bernard William et al
Wyeth Laboratories,
Patents & Trade Marks Department,
Huntercombe Lane South,
Taplow
Maidenhead, Berkshire SL6 0PH (GB)**

(54) Fungicidal mixtures


(57) The invention relates to a novel fungicidal composition comprising a fungicidally acceptable carrier and/or surface active agent together with, and synergistically effective amounts of

(a) at least one benzophenone of formula I



wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷, m and n have the meaning given; at least one fungicidal active ingredient selected from the following groups (A), (B), (C), (D) and (E):

- (A) an ergosterol biosynthesis inhibitor;
- (B) a strobilurine derivative,
- (C) a melanin biosynthesis inhibitor;
- (D) a compound selected from the group consisting of acibenzolar, benomyl, captan, carboxin, chlorothalonil, copper, cyprodinil, dinocap, dithianon, dimethomorph, dodine, ethirimol, famoxadone, fenpiclonil, fluazinam, mancozeb, metalaxyl, pyrifenoxy, sulfur, vinclozolin and
- (E) an azolopyrimidine of formula II



in which R^8 , R^9 , R^{10} , R^{11} , A, L and p have the meaning given;


and to a method of controlling the growth of phytopathogenic fungi at a locus which comprises applying synergistically effective amounts of (a) at least one benzophenones of formula I and (b) at least one fungicidal active ingredient selected from the groups (A), (B), (C), (D) and (E) to the locus.

Description

BACKGROUND OF THE INVENTION

[0001] The present invention relates to a fungicidal composition comprising a fungicidally acceptable carrier and/or surface active agent and synergistically effective amounts of

(a) at least one benzophenone of formula I




wherein

- R¹ represents a halogen atom, an optionally substituted alkyl, alkanoyloxy or alkoxy group; or a hydroxy group;
 R² represents a halogen atom or an optionally substituted alkyl group;
 m is 0 or an integer of 1 to 3;
 R³ independently represents a halogen atom, an optionally substituted alkyl or alkoxy group or a nitro group;
 R⁴ represents a halogen atom, a cyano, carboxy, hydroxy or nitro group or an optionally substituted alkyl, alkoxy, alkenyl, alkylthio, alkylsulphinyl, alkylsulphonyl or amino group;
 R⁵ represents an optionally substituted alkyl group;
 R⁶ represents a halogen atom or a nitro group, an optionally substituted alkyl, alkoxy, alkenyloxy, alkynyloxy, alkylthio, cycloalkyl, cycloalkyloxy, aryloxy group;
 n is 0 or 1; and
 R⁷ independently represents a halogen atom, an optionally substituted alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy, cycloalkyl, cycloalkoxy group;

(b) at least one fungicidally active ingredient selected from the following groups (A), (B), (C), (D) and (E):

- (A) an ergosterol biosynthesis inhibitor;
 (B) a strobilurine derivative,
 (C) a melanin biosynthesis inhibitor,
 (D) a compound selected from the group consisting of acibenzolar, benomyl, captan, carboxin, chlorothalonil, copper, cyprodinil, dinocap, dithianon, dimethomorph, dodine, ethirimol, famoxadone, fenpiclonil, fluazinam, mancozeb, metalaxyl, pyrifenoxy, sulfur and vinclozolin, and
 (E) an azolopyrimidine of formula II



in which

R⁸ and R⁹ each independently represent hydrogen or an optionally substituted alkyl, alkenyl, alkynyl, alka-

- R^8 and R^9 dieny, aryl, heteroaryl, cycloalkyl, bicycloalkyl or heterocyclyl group, or
 R^{10} together with the interjacent nitrogen atom represent an optionally substituted heterocyclic ring,
 R^{11} represents hydrogen or an alkyl or aryl group,
 L represents a hydrogen or halogen atom or an alkyl or alkoxy group,
 A independently represents a halogen atom or an optionally substituted alkyl or alkoxy group,
 represents N or CR^{12} , wherein R^{12} has the meaning given for R^{10} , and p is 0 or an integer from 1 to 5.

[0002] The fungicidal compounds of formula I to be used according to the present invention are known for example from US patent US 5,773,663.

[0003] The compounds of the classes (A), (B) and (D) are known from The Pesticide Manual 11th edition 1997, Editor Clive Tomlin.

[0004] The class of melanin biosynthesis inhibitors (MBI) (C) are chemical compounds which are capable of diminishing the in-vivo synthesis of melanin by inhibiting any of the reductase and/or dehydratase enzymes which are responsible for converting tetrahydroxynaphthalene into dihydroxynaphthalene. This class of compounds includes the following known compounds: carpropamid, chlorthalozone, diclocymet, pyroquilon, phthalide, tricyclazole and certain phenoxyamides, which are known for example from EP 0 262 393, in particular AC 382042 and Japanese patent application JP 5-9165-A.

[0005] The fungicidal compounds of formula II to be used according to the present invention are known for example from US patent US 5,593,996 and from International patent applications WO 98/46607 and WO 98/46608.

[0006] US 5,773,663 suggests to combine fungicidal benzophenone derivatives with other fungicides such as 4,6-dinitro-o-cresol, benalaxyl, benomyl, captan, carbendazim, chlorothalonil, copper, cymoxanil, dichlofluanid, dichlone, difenoconazole, dimethomorph, diniconazole, dinocap, dithianon, fenpiclonil, fenpropiomorph, hymexazol, imazalil, iprodione, isoprothiolane, kasugamycin, mancozeb, mepronil, mercuric oxide, oxadixyl, oxolinic acid, penconazole, propineb, pyrifenoxy, thiabendazole, thiram, tolclofos-methyl, triadimefon, triflumizole, triforine, validamycin A, vinclozolin, zineb and ziram.

[0007] However, there is no hint that such mixtures show synergistic effects and can advantageously be used for controlling diseases such as wheat powdery mildew, wheat leaf rust and wheat Septoria leaf blotch, Botrytis diseases and others.

[0008] Surprisingly, a strong synergy between the compounds of formula I and the fungicidally active ingredients selected from the classes (A), (B), (C), (D) and (E) as described above in greenhouse and field trials was found when these two compounds were in-tank mixed and when the activity of these mixtures was compared with that of the solo activity of each active ingredient.

[0009] A mixture of fungicides shows synergistic effect if the fungicidal activity of the mixture is larger than the sum of activities of the separately applied compounds. The expected fungicidal activity for a given mixture of two fungicides can also be calculated as follows (See Colby, S.R., "Calculating synergistic and antagonistic response of herbicide combinations", Weeds 15, pp 20-22 (1967):

$$EE = x + y - x \cdot y / 100$$

wherein

x is the efficacy in % compared with an untreated control upon treatment with a fungicidal active ingredient A at a dose rate a;

y is the efficacy in % compared with an untreated control upon treatment with a fungicidal active ingredient B at a dose rate b;

EE is the expected efficacy with a combination of fungicidal active ingredients A and B at a dose of a + b, respectively.

[0010] If the actual efficacy (E) exceeds the expected (calculated) one (EE), the mixture displays a synergistic effect.

SUMMARY OF THE INVENTION

[0011]

(b) The present invention includes a fungicidal composition comprising an acceptable carrier and/or surface active agent and synergistically effective amounts of at least one compound of formula I, and at least one fungicidal active


ingredient selected from the following the following groups (A), (B), (C), (D) and (E):

- (A) an ergosterol biosynthesis inhibitor;
- (B) a strobilurine derivative,
- (C) a melanin biosynthesis inhibitor,
- (D) a compound selected from the group consisting of acibenzolar (BION), benomyl, captan, carboxin, chlorothalonil, copper, cyprodinil, dinocap, dithianon, dimethomorph, dodine, ethirimol, famoxadone, fenpiclonil, fluazinam, mancozeb, metalaxyl, pyrifenoxy, sulfur and vinclozolin, and
- (E) an azolopyrimidine of formula II.

[0012] The present invention also includes a method of controlling the growth of phytopathogenic fungi at a locus which comprises applying synergistically effective amounts of at least one benzophenone of formula I and at least one fungicidally active ingredient selected from the following classes (A), (B), (C), (D) and (E) as defined above to the locus.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0013] Preferred compounds of formula I are benzophenones of formula IA,



wherein

R¹ represents a halogen atom, a methyl, trifluoromethyl, methoxy or hydroxy group, in particular a chlorine atom, a methyl or methoxy group;

R² represents a halogen atom, in particular a chlorine atom or a methyl group;

R³ represents a bromine or chlorine atom, a methyl, trifluoromethyl or nitro group, in particular a bromine atom;

R⁴ represents a methyl group;

R⁵ represents an alkyl group, in particular a methyl group;


R⁶ and R⁷ each independently represent an alkoxy group which may be substituted by a phenyl, alkylphenyl or halophenyl group, preferably C1-6 alkoxy being optionally substituted by a phenyl, methylphenyl or fluorophenyl group, in particular methoxy, benzyloxy and 2-fluorobenzyloxy; and

n is 0 or 1.

[0014] Particularly preferred are the benzophenones selected from the group consisting of 6'-butoxy-2,6-dichloro-4',5'-dimethoxy-2'-methylbenzophenone coded BP-1, 2,6-dichloro-4',5'-dimethoxy-6'-(2-fluorobenzyloxy)-2'-methylbenzophenone coded BP-2, 6'-benzyloxy-4',5'-dimethoxy-2,6-dimethyl-2'-methylbenzophenone coded BP-3; 3-bromo-2',6-dimethyl-2,4',5',6'-tetramethoxybenzophenone coded BP-4 and 2,6-dichloro-2'-methyl-4',5',6'-trimethoxybenzophenone coded BP-5, most preferred is BP-4.


[0015] Preferred ergosterol biosynthesis inhibitors of group (A) are selected from the group consisting of fenarimol, fenpropimorph, fenpropidine, spiroxamine and triforine.

[0016] Another group of ergosterol biosynthesis inhibitors are azole derivatives of formulae IIIA and IIIB,




wherein

E represents a linking group selected from the groups (a), (b), (c), (d) and (e):



in which

- X¹ represents an alkyl or an optionally substituted phenyl group;
- X² and X³ each independently represent a hydrogen atom or an alkyl group;
- X⁴ represents an alkyl or cyclopropylalkyl group;
- X⁵ represents a hydroxy or cyano group;
- X⁶ represents an optionally substituted phenyl group;
- X⁷ represents a halogen atom;
- q is 1, 2 or 3; and
- r is 0 or 2;




wherein X⁷ and q have the meaning given for formula IIIA, and E represents a group of formula -N(X⁸)-(CH₂)_s-O-, in which X⁸ represents a hydrogen atom or an alkyl group and s is an integer from 1 to 6.

[0017] Particularly preferred azole derivatives of group (A) are selected from the group consisting of cyproconazole,

epoxiconazole, flusilazole, metconazole, myclobutanil, penconazole, prochloraz, propiconazole, tebuconazole, triadimefon and tridimenol. Most preferred are epoxiconazole, metconazole, myclobutanil and prochloraz.


[0018] Preferred strobilurine derivatives of group (B) are the compounds of formula IV,



wherein

W represents N or CH;

B represents a -O-, -OCH₂-, a -CH₂O-, a pyrimid-4,6-dioxydiyl group or a group of formula



R¹² represents a C₁₋₄ alkyl group;

R¹³ represents a C₁₋₆ alkoxy or a C₁₋₆ alkylamino group;


R¹⁴ represents a hydrogen or halogen atom or a cyano, a C₁₋₄ alkyl or a C₁₋₄ haloalkyl group; and

s is 0, 1 or 2; in particular azoxystrobin, kresoxim methyl, trifloxystrobin or SSF126, most preferred azoxystrobin and kresoxim methyl.

[0019] Preferred melanin inhibitors of group (C) are selected from the group consisting of capropamid, chiobenthiazone, diclocymet, pyroquilon, phthalide, tricyclazole and a phenoxamide coded AC 382042, most preferred is AC 382042.

[0020] Preferred compounds of group (D) are selected from the group consisting of acibenzolar (BION), cyprodinil, dodine, ethirimol, famoxadone, fenpiclonil, fluazinam, mancozeb and metalaxyl.

[0021] Preferred azolopyrimidines of group (E) are the compounds of formula IIA,



wherein

R⁸ represents a C₁₋₈ alkyl, C₂₋₈ alkenyl, C₃₋₈ cycloalkyl or C₁₋₈ haloalkyl group, in particular a straight-chained or branched C₃₋₆ alkyl, C₂₋₆ alkenyl, C₅₋₆ cycloalkyl or C₂₋₆ fluoroalkyl group, most preferred an isopropyl, 2-butyl, cyclopentyl, methallyl, 2,2,2-trifluoroethyl or 1,1,1-trifluoroprop-2-yl group; and

R⁹ represents a hydrogen atom, or a C₁₋₈ alkyl group; most preferred a hydrogen atom or a methyl or

ethyl group; or

R⁸ and R⁹

together form an optionally substituted alkylene group having 3 to 6 carbon atoms in the main chain, in which one CH₂ group may be replaced by O, S or NH, in particular a piperid-1-yl group being optionally substituted by a C₁₋₆ alkyl group, most preferred a 4-methylpiperid-1-yl group;

L¹, L² and L³

each independently represent a hydrogen or halogen atom or a C₁₋₄ alkoxy group, at least one of which represents a halogen atom, in particular wherein L¹ represents a fluorine atom, L² represents a hydrogen or fluorine atom or a methoxy group and L³ represents a fluorine or chlorine atom; and

Hal

denotes a halogen atom, in particular a chlorine atom.

Particularly preferred are the azolopyrimidines of formula II selected from the group consisting of 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(cyclopropylamino)-[1,2,4]triazolo[1,5-a]pyrimidine coded AP-1, 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(4-methylpiperid-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine coded AP-2, 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(2,2,2-trifluoroethylamino)-[1,2,4]triazolo[1,5-a]pyrimidine coded AP-3 and 5-chloro-6-(2,4,6-trifluorophenyl)-7-[2-(1,1,1-trifluoro)propylamino]-[1,2,4]triazolo[1,5-a]pyrimidine coded AP-4, most preferred is AP-4.

[0022] A particularly preferred embodiment of this invention are compositions of three active ingredients which comprise one compound of formula I and two different compounds selected from the groups (A), (B), (C), (D) and (E), preferably one strobilurine compound selected from group (B) and one ergosterol biosynthesis inhibitor of group (A), in particular kresoxim-methyl and epoxiconazole or kresoxim-methyl and fenpropimorph.

[0023] Another particularly preferred embodiment of the invention are compositions comprising one compound of formula I and two different compounds of group (D), in particular dimethomorph and mancozeb.

[0024] Preferred are co-formulations, comprising the following constituents:

- a carrier;
- at least one benzophenone of formula I,
- at least one compound selected from the classes (A) through (E) as defined above;
- optionally an adjuvant selected from the group consisting of polyalkoxylated alcohols, triglycerides and amines, in particular Synperonic 91-6, which is commercially available from Uniqema, formerly ICI Surfactants;
- optionally a foam breaking agent.

[0025] The compound of formula I and the compound selected from the classes (A) through (E) as defined above are to be applied together, in synergistically effective amounts. These synergistic mixtures exhibit an extraordinary efficacy against a broad range of phytopathogenic fungi, in particular against fungi from the classes ascomycetes, basidiomycetes, oomycetes and deuteromycetes. Therefore, they can be applied advantageously against a broad range of diseases in different crops. They may be applied as leaf, stem, root, into-water, seed dressing, nursery box or soil fungicides.

[0026] The mixture according to the invention may be preferably applied for controlling phytopathogenic fungi of the genera:

Achlya, *Altemaria*, *Balansia*, *Bipolaris*, *Blumeria*, *Botrytis*, *Cercospora*, *Cochliobolus*, *Curvularia*, *Cylindrocladium*, *Drechslera*, *Etiyoma*, *Erysiphe*, *Fusarium*, *Gaeumannomyces*, *Gerlachia*, *Gibberella*, *Guignardia*, *Leptosphaeria*, *Magnaporthe*, *Mucor*, *Mycosphaerella*, *Myrothecium*, *Nigrospora*, *Pemnospora*, *Phoma*, *Pseudoperonospora*, *Pseudocercospora*, *Phytophthora*, *Puccinia*, *Pyrenophora*, *Pyricularia*, *Pythium*, *Rhizoctonia*, *Rhizopus*, *Rhynchosporium*, *Sarocladium*, *Sclerophthora*, *Sclerotium*, *Septoria*, *Tilletia*, *Uncinula*, *Ustilago*, *Ustilagoidea*, and *Venturia*, in particular the species *Blumeria graminis* f. sp. *tritici*, *Cercospora beticola*, *Septoria tritici*, *Erysiphe cichoracearum*, *Puccinia recondita* and *Pyrenophora teres*.

The mixtures according to the invention are in particular applied for controlling the above phytopathogenic fungi on monocotyledonous plants, such as barley and wheat, rice and turf grasses or fruit crops such as pomefruits, stonefruits and vines as well as all kinds of vegetables and ornamentals.

[0027] The application rate of the compound of formula I according to this invention is usually in the range of 1 to 2000 grams of active ingredient (g a.i.) per hectare, with rates between 20-500 g a.i./ha often achieving satisfactory control. The optimal rate for a specific application will depend on the crop(s) under cultivation and the predominant species of infesting fungi, and readily may be determined by established biological tests known to those skilled in the art.

[0028] In general, the preferred application rate of the compounds of formula I is in the range of 10 to 500 g a.i./ha, preferably 20-300 g a.i./ha.

[0029] The optimal rate for the compound of group (b) including the classes (A) through (E) will, however, depend on the crop(s) under cultivation and the level of infestation by the fungus, and can readily be determined by established biological tests.

[0030] The ratio (by weight) of the compound of formula I to the fungicidal active ingredient of the classes (A) through (E) as defined above is as a rule, from 100: 1 to 1: 100. The preferred ratio formula I : (A) through (E) may vary, e.g., from about 10: 1 to about 1 : 10, in particular from about 5 : 1 to about 1 : 5, most preferred from 2 : 1 to 1 : 2.

[0031] In the three-ways-compositions according to the present invention, i.e. the compositions containing one compound of formula I and two different compounds selected from the classes (A) through (E), the preferred relative ratios (by weight) are as follows:

compound of formula I: 200 to 1, preferably 20 to 1

1st compound of (A) to (E): 1 to 100, preferably 1 to 10

2nd compound of (A) to (E): 1 to 100, preferably 1 to 10.

[0032] The active compounds can be co-formulated together in a suitable ratio according to the present invention, together with usual carriers or diluents and/or additives known in the art.

[0033] Accordingly the invention further provides a fungicidal composition which comprises a carrier and, as active ingredient, at least one compound of formula I as defined above and at least one fungicidal active ingredient selected from the classes (A) through (E) as defined above.

[0034] A method of making such a composition is also provided which comprises bringing the compound of formula I and the fungicidal active ingredient selected from the classes (A) through (E) as defined above into association with at least one carrier. It is also envisaged that different isomers or mixtures of isomers of formula I and/or the fungicidal active ingredient selected from the classes (A) through (E) may have different levels or spectra of activity and thus compositions may comprise individual isomers or mixtures of isomers.

[0035] A composition according to the invention preferably contains from 0.1% to 99.9%, preferably 0.2 to 80 % by weight (w/w) of active ingredients.

[0036] A carrier in a composition according to the invention is any material with which the active ingredient is formulated to facilitate application to the locus to be treated, which may for example be a plant, seed, foliage, soil, or into the water where the plant grow, or to the roots or to facilitate storage, transport or handling. A carrier may be a solid or a liquid, including material which is normally a gas but which has been compressed to form a liquid.

[0037] The compositions may be manufactured into e.g. emulsion concentrates, solutions, oil in water emulsions, wettable powders, soluble powders, suspension concentrates, dusts, granules, water dispersible granules, tablets, micro-capsules, gels and other formulation types by well-established procedures. These procedures include intensive mixing and/or milling of the active ingredients with other substances, such as fillers, solvents, solid carriers, surface active compounds (surfactants), and optionally solid and/or liquid auxiliaries and/or adjuvants. The form of application such as spraying, atomizing, dispersing or pouring may be chosen like the compositions according to the desired objectives and the given circumstances.

[0038] Solvents may be aromatic hydrocarbons, e.g. Solvesso® 200, substituted naphthalenes, phthalic acid esters, such as dibutyl or dioctyl phthalate, aliphatic hydrocarbons, e.g. cyclohexane or paraffins, alcohols and glycols as well as their ethers and esters, e.g. ethanol, ethyleneglycol mono- and dimethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, or γ -butyrolactone, higher N-alkylpyrrolidones, e.g. N-octylpyrrolidone or N-cyclohexylpyrrolidone, epoxidized plant oil esters, e.g. methylated coconut or soybean oil ester and water. Mixtures of different liquids are often suitable.

[0039] Solid carriers, which may be used for dusts, wettable powders, water dispersible granules, or granules, may be mineral fillers, such as calcite, talc, kaolin, montmorillonite or attapulgite or others. The physical properties may be improved by addition of highly dispersed silica gel or polymers. Carriers for granules may be porous material, e.g. pumice, kaolin, sepiolite, bentonite; non-sorptive carriers may be calcite or sand or others. Additionally, a multitude of pregranulated inorganic or organic materials may be used, such as dolomite or crushed plant residues.

[0040] Pesticidal compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier which is a surfactant facilitates this process of dilution. Thus, preferably at least one carrier in a composition according to the invention is a surfactant. For example, the composition may contain at two or more carriers, at least one of which is a surfactant.

[0041] Surfactants may be nonionic, anionic, cationic or zwitterionic substances with good dispersing, emulsifying and wetting properties depending on the nature of the compound according to general formula I to be formulated. Surfactants may also mean mixtures of individual surfactants.

[0042] The compositions of the invention may for example be formulated as wettable powders, water dispersible granules, dusts, granules, solutions, emulsifiable concentrates, emulsions, suspension concentrates and aerosols. Wettable powders usually contain 5 to 90% w/w of active ingredient and usually contain in addition to solid inert carrier, 3 to 10% w/w of dispersing and wetting agents and, where necessary, 0 to 10% w/w of stabilizer(s) and/or other additives such as penetrants or stickers. Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and may be diluted in the field with further solid carrier to give a com-

position usually containing 0.5 to 10% w/w of active ingredient. Water dispersible granules and granules are usually prepared to have a size between 0.15 mm and 2.0 mm and may be manufactured by a variety of techniques. Generally, these types of granules will contain 0.5 to 90% w/w active ingredient and 0 to 20% w/w of additives such as stabilizer, surfactants, slow release modifiers and binding agents. The so-called "dry flowables" consist of relatively small granules having a relatively high concentration of active ingredient. Emulsifiable concentrates usually contain, in addition to a solvent or a mixture of solvents, 1 to 80% w/v active ingredient, 2 to 20% w/v emulsifiers and 0 to 20% w/v of other additives such as stabilizers, penetrants and corrosion inhibitors. Suspension concentrates are usually milled so as to obtain a stable, non-sedimenting flowable product and usually contain 5 to 75% w/v active ingredient, 0.5 to 15% w/v of dispersing agents, 0.1 to 10% w/v of suspending agents such as protective colloids and thixotropic agents, 0 to 10% w/v of other additives such as defoamers, corrosion inhibitors, stabilizers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation and crystallization or as antifreeze agents for water.

[0043] Aqueous dispersions and emulsions, for example compositions obtained by diluting the formulated product according to the invention with water, also lie within the scope of the invention.

[0044] Of particular interest in enhancing the duration of the protective activity of the compounds of this invention is the use of a carrier which will provide slow release of the pesticidal compounds into the environment of a plant which is to be protected.

[0045] The biological activity of the active ingredient can also be increased by including an adjuvant in the formulation or the spray dilution. An adjuvant is defined here as a substance which can increase the biological activity of an active ingredient but is not itself significantly biologically active. The adjuvant can either be included in the formulation as a coformulant or carrier, or can be added to the spray tank together with the formulation containing the active ingredient.

[0046] As a commodity the compositions may preferably be in a concentrated form whereas the end user generally employs diluted compositions. The compositions may be diluted to a concentration down to 0.001% of active ingredient. The doses usually are in the range from 0.01 to 10 kg a.i./ha.

[0047] Examples of formulations which can be used according to the invention are:

SC-I 1

active ingredient	<u>BP-1</u>	100.0 g
Dispersing agent	Morwet D425 ¹⁾	25.0 g
Dispersing agent	Pluronic® PE10500 ²⁾	5.0 g
Antifoaming agent	Rhodorsil® 426R ³⁾	1.5 g
Dispersing agent	Rhodopol® 23 ³⁾	2.0 g
Antifreezing agent	Propylene glycol	80.0 g
Biocidal agent	Proxel® GXL ⁴⁾	1.0 g
Water		to 1000 ml

SC-I 2

active ingredient	<u>BP-4</u>	100.0 g
Dispersing agent	Soprophor® FL ³⁾	30.0 g
Antifoaming agent	Rhodorsil® 426R ³⁾	1.5 g

Dispersing agent	Rhodopol® 23 ³⁾	2.0 g
Antifreezing agent	Propylene glycol	80.0 g
Biocidal agent	Proxel® GXL ⁴⁾	1.0 g
Water		to 1000 ml

SC-A-E

active ingredient	fungicide selected from classes (A) through (E)	200.0 g
Dispersing agent	Soprophor® FL ³⁾	25.0 g
Antifoaming agent	Rhodorsil® 426R ³⁾	1.5 g
Dispersing agent	Rhodopol® 23 ³⁾	2.0 g
Antifreezing agent	Propylene glycol	80.0 g
Biocidal agent	Proxel® GXL ⁴⁾	1.0 g
Water		to 1000 ml

SC-I + A-E

active ingredient	<u>BP-4</u>	60.0 g
active ingredient	fungicide selected from classes (A) through (E)	120.0 g
Dispersing agent	Soprophor® FL ³⁾	25.0 g
Antifoaming agent	Rhodorsil® 426R ³⁾	1.5 g
Dispersing agent	Rhodopol® 23 ³⁾	2.0 g
Antifreezing agent	Propylene glycol	80.0 g
Biocidal agent	Proxel® GXL ⁴⁾	1.0 g
Water		to 1000 ml

DC-I 1

active ingredient	<u>BP-4</u>	100.0 g
Wetting agent	Pluronic® PE6400 ²⁾	50.0 g
Dispersing agent	Lutensol® TO 12 ²⁾	50.0 g
Solvent	benzyl alcohol	to 1000 ml

- 1) Product commercially available from Witco
- 2) Product commercially available from BASF AG, Germany
- 3) Product commercially available from Rhône-Poulenc
- 4) Product commercially available from Zeneca

[0048] The formulation SC-A-E comprising a compound selected from the classes (A) through (E) is in-tank mixed with any of the other formulations SC-I 1, SC-I 2, SC-I 3, or DC-I which comprise the compound of formula I.

[0049] In a preferred embodiment the active ingredients are added to the tank mix together each as solo formulation.

[0050] Therefore, the present invention relates to a kit for the preparation of a spray mixture consisting of two separate containments:

(i) a containment which comprises at least one benzophenone of formula I, in particular one or more compounds selected from BP-1 through BP-4 conventional carriers and optionally adjuvants;

(ii) a containment which comprises at least one active ingredient selected from the classes (A) through (E).

[0051] In a preferred embodiment the said kit will consist of two bottles with dispensing means which allow the easy and correct addition of the active ingredients (a) and (b) to the tank mix.

[0052] The formulation SC-I + A-E comprising BP-4 and a fungicidal active ingredient selected from the classes (A) through (E) as defined above can be used directly for preparing the tank mix according to the present invention.

[0053] A composition according to the invention preferably contains from 0.5% to 95% by weight of active ingredients.

[0054] As commodity the compositions may preferably be in a concentrated form whereas the end-user generally employs diluted compositions. The compositions may be diluted down to a concentration of 0.0001% of active ingredients.

[0055] The compositions of this invention can be applied to the plants or their environment simultaneous with or in succession with other active substances. These other active substances can be either fertilizers, agents which donate trace elements or other preparations which influence plant growth. However, they can also be other fungicides, selective herbicides, insecticides, bactericides, nematocides, algicides, molluscicides, rodenticides, virucides, compounds inducing resistance into plants, biological control agents such as viruses, bacteria, nematodes, fungi and other microorganisms, repellents of birds and animals, and plant growth regulators, or mixtures of several of these preparations, if appropriate together with other carrier substances conventionally used in the art of formulation, surfactants or other additives which promote application.

[0056] Examples of insecticidal compounds are alpha-cypermethrin, benfuracarb, BPMC, buprofezine, carbosulfan, cartap, chlorfenvinphos, chlorpyrifos-methyl, cycloprothrin, cypermethrin, esfenvalerate, ethofenprox, fenpropathrin, flucythrinate, flufenoxuron, hydramethylnon, imidacloprid, isoxathion, MEP, MPP, nitenpyram, PAP, permethrin, propaphos, pymetrozine, silafluofen, tebufenozide, teflubenzuron, temephos, terbufos, tetrachlorvinphos and triazamate.

[0057] Examples of biological control agents are: *Bacillus thuringiensis*, *Verticillium lecanii*, *Autographa californica* NPV, *Beauveria bassiana*, *Ampelomyces quisqualis*, *Bacillus subtilis*, *Pseudomonas chlororaphis*, *Pseudomonas fluorescens*, *Streptomyces griseoviridis* and *Trichoderma harzianum*.

[0058] Examples of chemical agents that induce systemic acquired resistance in plants such as: isonicotinic acid or derivatives thereof, 2,2-dichloro-3,3-dimethylcyclopropylcarboxylic acid.

[0059] The present invention is of wide applicability in the protection of crops, trees, residential and ornamental plants against fungal attack. Preferred crops are cereals, such as wheat and barley, rice as well as vines and apples. The duration of the protection is normally dependent on the individual compound selected, and also a variety of external factors, such as climate, whose impact is normally mitigated by the use of a suitable formulation.

[0060] The following examples further illustrate the present invention. It should be understood, however, that the invention is not limited solely to the particular examples given below.

EXAMPLES

General Methods

[0061] The trials are carried out under greenhouse (Examples 1 to 18) or field conditions (Example 19) in residual or curative applications. The fungicides are applied in single treatments, or in a combination comprising a benzophenone of formula I and a compound selected from the classes (A) through (E) as defined above. The compounds are applied in form of an aqueous spray mix obtained from concentrated formulation or the technical material.

I. Cereals and Dicots - Greenhouse

[0062]

1. Seed is planted in 6 cm diameter plastic pots and maintained in the greenhouse.
2. When the primary leaf is fully expanded in the case of cereals or several leaves are present in the case of dicots, formulated test compounds are sprayed with a three nozzle overhead fungicide sprayer to near run-off. Alternatively, a single nozzle overhead track sprayer is used for application of the compounds to cereals at a rate of 200 l/ha.

Plants are then allowed to air-dry.

3. Inoculation precedes treatment in the case of curative evaluations and follows treatment in case of residual evaluations.

For inoculation of powdery mildew disease, plants are set up on greenhouse benches with bottom watering mats and inoculated by dusting them with conidia from infected plants. Between inoculation and treatment for curative evaluations and between treatment and inoculation for residual evaluations, plants are maintained in the greenhouse with bottom watering.

For inoculation of non-powdery mildew diseases, an aqueous spore suspension of the pathogen is applied to the plant and the plants are kept 1-2 days in a moist infection chamber before being returned to the greenhouse where they are maintained by bottom watering.

4. Disease on the foliage as percent leaf area with disease symptoms/signs is evaluated about 7 days after inoculation. In the case of wheat, the tips and bases of the leaves are excluded from the evaluation.

$$\% \text{ disease control} = 100 - \frac{\% \text{ disease in treated plants}}{\% \text{ disease in untreated plants}} \times 100\%$$

FORMULATION, REFERENCE COMPOUNDS AND CONTROLS:

[0063]

1. Technical compounds are formulated in a solvent/surfactant system consisting of 5% acetone and 0.05% Tween

20 in deionized water.

Compounds are dissolved in acetone prior to addition of the water; the Tween 20 can be added through either the acetone or the water. Dilutions are made using the solvent/surfactant system.

Formulated compounds are prepared using deionized water.

2. Two kinds of controls are included:

Plants treated with the solvent/surfactant solution and inoculated (Solvent Blank).

[0064] Untreated plants which are inoculated (Inoculated Control). For the field study formulated benzophenones BP-1 through BP-4 and formulated compounds from the classes (A) through (E) were used.

Evaluation of the disease:

[0065] Assessments of the diseases took place at the indicated day after the application of the compounds. Percent infected leaf area infected was evaluated. The efficacy of the compounds/compounds mixtures to control the diseases was calculated by using the formula given above under item 4:

II. Apple Fruit Botrytis

[0066]

1. Apples (*Malus X domestica* Borkh.) variety "Golden Delicious" are disinfected by washing them briefly in 70 % ethanol. After drying the apples are marked with four short equal-distant lines indicating the positions to be wounded.

2. Corresponding with the marks, four holes are poked around the apple equator with a pipette tip. 10 µl of the treatment solution are pipetted into each hole.

3. Three hours after application, 10 µl of a conidial suspension of *Botrytis cinerea* are pipetted into each hole. For incubation, the treated/inoculated apples are stored for five days.

4. Disease occurs as rotten apple tissue surrounding the inoculated wounds. The diameter of the rotten zone around each wound is measured.

FORMULATION, REFERENCE COMPOUNDS AND CONTROLS:1.

Technical compounds are formulated in a solvent system consisting of 5% acetone and 0.05% Tween 20 in deionized water. Compounds are dissolved in acetone prior to dilution with water. Formulated compounds are prepared using deionized water.

2. Three kinds of controls are included:

Apples treated with the solvent solution and inoculated (Solvent Blank).

Untreated apples which are inoculated (Inoculated Control).

Untreated apples which are not inoculated (Uninoculated Control).

Evaluation of the disease:

[0067] Assessments of the diseases took place at the indicated day after the application of the compounds. Percent infected leaf area infected was evaluated. The efficacy of the compounds/compounds mixtures to control the diseases was calculated by using the formula:

$$\% \text{ disease control} = 100 - \frac{\text{mean of diameters on treated apples}}{\text{mean of diameters on untreated apples}} \times 100\%$$

Determination of synergy:

[0068] Synergy was calculated using the % disease control values of specific treatments for the two COLBY formula given hereinabove

III. Field Tests

[0069] The compounds are applied according to good agricultural practice in form of an aqueous spray mix obtained from concentrated formulation or the technical material at a rate of 400 l/ha. The disease control is evaluated according to the formula given for the greenhouse tests.

A Greenhouse TestsExample 1

Fungicidal efficacy of the mixture of BP-1 + AP-1 (4 day curative) against *Erysiphe graminis* on wheat

[0070] The tank mix was obtained from technical materials of BP-1 and AP-1. The observed and expected efficacies with different rates are given in Table I:

Table I

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-1</u>	<u>AP-1</u>		
125	0	42	--
25	0	1	--
0	125	0	--
0	25	0	--
125	125	56	42
125	25	54	42
25	125	21	1
25	25	4	1

Example 2

Fungicidal efficacy of the mixture of BP-1 + AP-2 (4 day curative) against *Erysiphe graminis* on wheat

[0071] The tank mix was obtained from technical materials of BP-1 and AP-2. The observed and expected efficacies with different rates are given in Table II:

Table II

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-1</u>	<u>AP-2</u>		
125	0	42	--
25	0	1	--
0	125	8	--
0	25	0	--
125	125	67	47

EP 1 023 834 A1

Table II (continued)

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-1</u>	<u>AP-2</u>		
125	25	73	42
25	125	20	9
25	25	9	1

Example 3

Fungicidal efficacy of the mixture of BP-1 + triadimefon (4 day curative) against *Erysiphe graminis* on wheat

[0072] The tank mix was obtained from technical material of BP-1 and a wettable powder formulation containing 250 g/kg triadimefon. The observed and expected efficacies with different rates are given in Table III:

Table III

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-1</u>	triadimefon		
125	0	42	--
25	0	1	--
0	125	30	--
0	25	12	--
125	125	85	59
125	25	56	49
25	125	41	31
25	25	6	13

Example 4

Fungicidal efficacy of the mixture of BP-5 + triforine (3 day protective) against *Erysiphe graminis* on wheat

[0073] The tank mix was obtained from technical material of BP-5 and an EC formulation containing 190 g/l triforine. The observed and expected efficacies with different rates are given in Table IV:

Table IV

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-5</u>	triforine		
25	0	31	--
5	0	12	--
0	125	12	--
25	125	59	41
5	25	26	24

Example 5

Fungicidal efficacy of the mixture of BP-1 + triadimefon (3 day protective) against *Erysiphe graminis* on wheat

[0074] The tank mix was obtained from technical material of BP-1 and a wettable powder formulation containing 250 g/kg triadimefon. The observed and expected efficacies with different rates are given in Table V:

Table V

dose rate (ppm)		Observed Efficacy	Expected Efficacy
<u>BP-1</u>	triadimefon		
5	0	78	--
1	0	44	
0	25	24	--
5	25	90	83
1	25	78	57

Example 6

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Erysiphe cichoracearum* on cucumbers

[0075] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table VI:

Table VI

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	64	100	
		16	51	
		4	6	
		1	1	
Dithianon	WG 700 g/kg	256	0	
		64	0	
		16	0	
		4	0	
Cyprodinil	TC 100 %	256	15	
		64	0	
		16	0	
		4	0	
Triforine	EC 190 g/l	256	96	
		64	79	
		16	44	
		4	1	
Fenpropidin	EC 750 g/l	256	69	
		64	21	
		16	6	
		4	0	
Mancozeb	WP 800 g/kg	256	33	
		64	1	
		16	0	

		4	0	
5	Quinoxifen TC 100 %	256	100	
		64	100	
		16	100	
		4	90	
10	Chlorothalonil SC 500 g/l	256	1	
		64	0	
		16	0	
		4	0	
15	Ethirimol SC 280 g/l	256	100	
		64	94	
		16	83	
		4	50	
20	Dimethomorph TC 100 %	256	18	
		64	0	
		16	0	
		4	0	
25	BION WG 500 g/kg	256	0	
		64	0	
		16	0	
		4	0	
30	Azoxystrobin TC 100 %	256	100	
		64	100	
		16	96	
		4	78	
35	BP-4 + Tankmix	64 + 256	100	100
	Dithianon	16 + 64	73	51
40		4 + 16	31	6
		1 + 4	29	1
	BP-4 + Tankmix	64 + 256	92	100
45	Cyprodinil	16 + 64	59	51
		4 + 16	51	6
		1 + 4	32	1
50	BP-4 + Tankmix	64 + 256	100	100
	Triforine	16 + 64	99	90
		4 + 16	65	47
5		1 + 4	31	3

5	BP-4 + Fenpropidin	Tankmix	64 + 256	100	100
			16 + 64	96	61
			4 + 16	60	12
			1 + 4	56	1
10	BP-4 + Mancozeb	Tankmix	64 + 256	100	100
			16 + 64	99	52
			4 + 16	54	6
			1 + 4	38	1
15	BP-4 + Quinoxifen	Tankmix	64 + 256	100	100
			16 + 64	100	100
			4 + 16	100	100
			1 + 4	100	91
20	BP-4 + Chlorothalonil	Tankmix	64 + 256	100	100
			16 + 64	99	51
			4 + 16	22	6
			1 + 4	14	1
25	BP-4 + Ethirimol	Tankmix	64 + 256	100	100
			16 + 64	100	97
			4 + 16	85	84
			1 + 4	53	51
30	BP-4 + Dimethomorph	Tankmix	64 + 256	100	100
			16 + 64	99	51
			4 + 16	37	6
			1 + 4	10	1
35	BP-4 + BION	Tankmix	64 + 256	100	100
			16 + 64	53	51
			4 + 16	12	6
			1 + 4	1	1
40	BP-4 + Azoxystrobin	Tankmix	64 + 256	100	100
			16 + 64	100	100
			4 + 16	99	96
			1 + 4	92	78

Example 7

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Puccinia recondite* on wheat

[0076] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table VII:

Table VII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	256	13	
		64	0	
		16	0	
		4	0	
Dithianon	WG 700 g/kg	256	95	
		64	84	
		16	16	
		4	4	
Cyprodinil	TC 100 %	256	0	
		64	0	
		16	0	
		4	0	
Triforine	EC 190 g/l	256	100	
		64	90	
		16	7	
		4	0	
Fenpropidin	EC 750 g/l	256	97	
		64	8	
		16	0	
		4	0	
Mancozeb	WP 800 g/kg	256	91	
		64	28	
		16	0	
		4	0	

5	Quinoxifen	TC 100 %	256	0	
			64	0	
			16	0	
			4	0	
10	Chlorothalonil	SC 500 g/l	256	31	
			64	0	
			16	0	
			4	0	
15	Ethirimol	SC 280 g/l	256	0	
			64	0	
			16	0	
20	Dimethomorph	TC 100 %	256	0	
			64	0	
			16	0	
25	BION	WG 500 g/kg	256	0	
			64	0	
			16	0	
			4	0	
30	BP-4 + Dithianon	Tankmix	256 + 256	100	95
			64 + 64	100	84
			16 + 16	100	16
			4 + 4	100	4
35	BP-4 + Cyprodinil	Tankmix	256 + 256	100	13
			64 + 64	100	0
			16 + 16	100	0
			4 + 4	100	0
40	BP-4 + Triforine	Tankmix	256 + 256	100	100
			64 + 64	100	90
			16 + 16	100	7
			4 + 4	100	0
45	BP-4 + Fenpropidin	Tankmix	256 + 256	100	97
			64 + 64	96	8
			16 + 16	93	0
			4 + 4	91	0
50	BP-4 + Mancozeb	Tankmix	256 + 256	100	92
			64 + 64	97	28

		16 + 16	94	0
		4 + 4	92	0
BP-4 +	Tankmix	256 + 256	89	13
Quinoxifen		64 + 64	96	0
		16 + 16	95	0
		4 + 4	93	0
BP-4 +	Tankmix	256 + 256	100	40
Chlorothalonil		64 + 64	36	0
		16 + 16	40	0
		4 + 4	9	0
BP-4 +	Tankmix	256 + 256	96	13
Ethirimol		64 + 64	96	0
		16 + 16	64	0
BP-4 +	Tankmix	256 + 256	54	13
Dimethomorph		64 + 64	28	0
		16 + 16	6	0
BP-4 +	Tankmix	256 + 256	75	13
BION		64 + 64	78	0
		16 + 16	64	0
		4 + 4	59	0

Example 8

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Leptosphaeria nodorum* on wheat

[0077] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table VIII:

Table VIII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	256	34	
		64	14	
		16	0	
		4	0	
Dithianon	WG 700 g/kg	256	74	
		64	37	
		16	1	
		4	0	
Cyprodinil	TC 100 %	256	93	
		64	88	
		16	75	
		4	0	
Triforine	EC 190 g/l	256	85	
		64	51	
		16	31	
		4	4	
		1	0	
Fenpropidin	EC 750 g/l	256	21	
		64	0	
Mancozeb	WP 800 g/kg	256	69	
		64	47	
		16	21	
		4	0	
Quinoxifen	TC 100 %	256	18	
		64	0	
		16	1	
Dimethomorph	TC 100 %	256	14	
		64	11	
		16	6	
		4	1	

5	Azoxystrobin	TC 100 %	256	100	
			64	100	
			16	97	
			4	84	
10	BP-4 + Dithianon	Tankmix	256 + 256	90	83
			64 + 64	65	46
			16 + 16	27	1
			4 + 4	4	0
15	BP-4 + Cyprodinil	Tankmix	256 + 256	97	96
			64 + 64	92	90
			16 + 16	80	75
			4 + 4	57	0
20	BP-4 + Triforine	Tankmix	256 + 256	97	90
			64 + 64	84	58
			16 + 16	47	31
			4 + 4	32	4
25	BP-4 + Fenpropidin	Tankmix	256 + 256	42	48
			64 + 64	21	14
30	BP-4 + Mancozeb	Tankmix	256 + 256	87	79
			64 + 64	65	55
			16 + 16	32	21
			4 + 4	8	0
35	BP-4 + Quinoxifen	Tankmix	256 + 256	62	46
			64 + 64	34	14
			16 + 16	8	1
40	BP-4 + CL 336 370 Dimethomorph	Tankmix	256 + 256	84	43
			64 + 64	65	24
			16 + 16	31	6
			4 + 4	14	1
45	BP-4 + Azoxystrobin	Tankmix	256 + 256	100	100
			64 + 64	100	100
			16 + 16	98	97
			4 + 4	95	84

Example 9

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Puccinia recondita* on wheat

[0078] The tank mixes were obtained from technical material of BP-4 and different formulations of different active

EP 1 023 834 A1

ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table IX:

Table IX

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	256	38	
		64	16	
		16	4	
		4	6	
Captan	WP 500 g/kg	256	89	
		64	55	
		16	11	
		4	4	
Fluazinam	SC 500 g/l	256	80	
		64	22	
		16	6	
		4	0	
Metalaxyl	TC 100 %	256	12	
		64	0	
Fenpiclonil	TC 100 %	256	58	
		64	18	
		16	0	
		4	1	
Famoxadone	TC 100 %	64	92	
		16	80	
		4	45	
BP-4 + Captan	Tankmix	256 + 256	95	93
		64 + 64	65	62
		16 + 16	30	15
BP-4 + Fluazinam	Tankmix	256 + 256	97	88
		64 + 64	53	35
		16 + 16	20	10
BP-4 + Metalaxyl	Tankmix	256 + 256	61	46
		64 + 64	19	16
BP-4+ Fenpiclonil	Tankmix	256 + 256	81	74
		64 + 64	54	31
		16 + 16	12	4
		4 + 4	7	7

EP 1 023 834 A1

Table IX (continued)

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4 + Famoxadone	Tankmix	64 + 64	93	93
		16 + 16	83	81
		4 + 4	60	48

Example 10

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Leptosphaeria nodorum* on wheat

[0079] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table X:

Table X

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	256	44	
		64	18	
		16	10	
		4	0	
Dodine	WP 650 g/kg	256	59	
		64	24	
		16	6	
		4	0	
Captan	WP 500 g/kg	256	90	
		64	84	
Fluazinam	SC 500 g/l	256	91	
		64	81	
Famoxadone	TC 100 %	256	97	
		64	80	
		16	69	
		4	66	
BP-4 + Dodine	Tankmix	256 + 256	75	77
		64 + 64	57	38
		16 + 16	29	15
		4 + 4	10	0
BP-4 + Captan	Tankmix	256 + 256	96	94
		64 + 64	93	87
BP-4 + Fluazinam	Tankmix	256 + 256	95	95
		64 + 64	91	84

EP 1 023 834 A1

Table X (continued)

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4 + Famoxadone	Tankmix	256 + 256	94	98
		64 + 64	89	84
		16 + 16	86	72
		4 + 4	67	66

Example 11

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Erysiphe cichoracearum* on cucumbers

[0080] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XI:

Table XI

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	50	96	
		10	55	
		2	2	
		0.4	0	
Sulfur inorganic	TC 100 %	50	0	
		10	7	
		2	0	
		0.4	0	
Propiconazole	EC 250 g/l	50	96	
		10	76	
		2	19	
		0.4	0	
Epoxiconazole	SC 125 g/l	50	77	
		10	68	
		2	46	
		0.4	0	
Tebuconazole	EC 250 g/l	50	95	
		10	79	
		2	45	
		0.4	10	
Metconazole	SL 60 g/l	50	99	
		10	75	
		0.4	2	
Myclobutanil	WP 60 g/kg	50	96	
		10	73	
		2	52	
		0.4	36	
Kresoxim- methyl	WG 500 g/kg	50	100	
		10	88	
		2	14	
		0.4	2	
AC 382042	TC 100 %	50	25	
		10	14	

		2	3
		0.4	7
5	BRIO SE 450 g/l	50	96
	Epoxiconazole / Fenpropimorph	10	48
	(150 g/l) / (300 g/l)		
10	ACROBAT MZ WP 690 g/kg	50	57
	Dimethomorph / Mancozeb	10	30
	(90 g/kg) / (600 g/kg)	2	0
		0.4	0
15	JUWEL SC 250 g/l	50	100
	Kresoxim-methyl / Epoxiconazole	10	98
	(125 g/l) / (125 g/l)	2	71
20		0.4	37
	BP-4 Tankmix	50 + 50	92 96
	Sulfur inorganic	10 + 10	70 58
25		2 + 2	10 2
		0.4 + 0.4	9 0
	BP-4 Tankmix	50 + 50	100 100
	Propiconazole	10 + 10	100 89
30		2 + 2	61 21
		0.4 + 0.4	28 0
	BP-4 Tankmix	50 + 50	100 99
	Epoxiconazole	10 + 10	98 86
35		2 + 2	71 47
		0.4 + 0.4	41 0
	BP-4 Tankmix	50 + 50	100 100
40	Tebuconazole	10 + 10	98 91
		2 + 2	48 45
		0.4 + 0.4	28 10
	BP-4 Tankmix	50 + 50	100 100
45	Metconazole	10 + 10	97 89
		0.4 + 0.4	5 2
	BP-4 Tankmix	50 + 50	100 100
50	Myclobutanil	10 + 10	99 88
		2 + 2	75 52
		0.4 + 0.4	55 36

BP-4	Tankmix	50 + 50	100	100
Kresoxim		10 + 10	100	95
-methyl		2 + 2	37	15
		0.4 + 0.4	2	2
BP-4	Tankmix	50 + 50	100	97
AC 382042		10 + 10	87	62
		2 + 2	28	5
		0.4 + 0.4	25	7
BP-4	Tankmix	50 + 50	100	100
BRIO		10 + 10	85	77
BP-4	Tankmix	50 + 50	100	98
ACROBAT MZ		10 + 10	89	69
		2 + 2	45	2
		0.4 + 0.4	7	0
BP-4	Tankmix	50 + 50	100	100
JUWEL		10 + 10	98	99
		2 + 2	84	72
		0.4 + 0.4	50	37

Example 12

Fungicidal efficacy of the mixture of BP-4 + Myclobutanil (4 day residual) against *Puccinia recondita* on wheat

[0081] The tank mixes were obtained from technical material of BP-4 and Myclobutanil. The type of formulation, the observed and expected efficacies with different rates are given in Table XII:

Table XII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	50	31	
		10	10	
		2	7	
		0.4	2	
Myclobutanil	WP 60 g/kg	50	99	
		10	72	
		2	52	
		0.4	13	

EP 1 023 834 A1

Table XII (continued)

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4 Myclobutanil	Tankmix	50 + 50	99	99
		10 + 10	86	75
		2 + 2	65	55
		0.4 + 0.4	48	14

Example 13

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Pyrenophora teres* on barley

[0082] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XIII:

Table XIII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	50	58	
		10	34	
		2	0	
		0.4	0	
Sulfur inorganic	TC 100 %	10	0	
		2	0	
Copper oxychloride	WP 450 g/kg	50	14	
		10	14	
		2	0	
		0.4	0	
Propiconazole	EC 250 g/l	50	74	
		10	38	
		2	0	
		0.4	0	
Metconazole	SL 60 g/l	10	47	
		0.4	0	
Myclobutanil	WP 60 g/kg	50	82	

EP 1 023 834 A1

		10	40	
		2	21	
		0.4	0	
Kresoxim-	WG 500 g/kg	2	21	
methyl		0.4	0	
ACROBAT MZ	WP 690 g/kg	50	47	
Dimethomorph/		10	23	
Mancozeb		2	12	
		0.4	0	
JUWEL	SC 250 g/l	50	89	
Kresoxim-		10	62	
methyl /		2	45	
Epoxiconazole		0.4	5	
BP-4	Tankmix	10 + 10	49	34
Sulfur inorganic		2 + 2	0	0
BP-4	Tankmix	50 + 50	78	64
Copper		10 + 10	54	43
oxychloride		2 + 2	38	0
		0.4 + 0.4	0	0
BP-4	Tankmix	50 + 50	89	89
Propiconazole		10 + 10	78	59
		2 + 2	32	0
		0.4 + 0.4	0	0
BP-4	Tankmix	50 + 50	89	86
Tebuconazole		2 + 2	14	5
		0.4 + 0.4	0	0
BP-4	Tankmix	10 + 10	69	65
Metconazole		2 + 2	25	32
		0.4 + 0.4	18	0
BP-4	Tankmix	50 + 50	91	93
Myclobutanil		10 + 10	67	61
		2 + 2	40	21
		0.4 + 0.4	7	0
BP-4	Tankmix	2 + 2	67	21
Kresoxim- methyl		0.4 + 0.4	36	0
BP-4	Tankmix	50 + 50	85	78
ACROBAT MZ		10 + 10	60	49

		2 + 2	29	12
		0.4 + 0.4	0	0
BP-4	Tankmix	50 + 50	100	95
JUWEL		10 + 10	78	75
		2 + 2	49	45
		0.4 + 0.4	40	5

15 Example 14

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Erysiphe cichoracearum* on cucumbers

20 [0083] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XIV:

25 Table XIV

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
30 BP-4	TC 100 %	50	97	
		10	13	
		2	0	
		0.4	0	
35 Cyproconazole	SL 100 g/l	10	68	
		0.4	19	
40 Dinocap	WP 190 g/kg	10	0	
		2	0	
		0.4	0	
45 Fenarimol	SC 125 g/l	50	67	
		10	42	
		2	34	
		0.4	5	
50 Fenpropimorph	EC 750 g/l	10	7	
		2	0	
		0.4	0	

Flusilazole	WP 200 g/kg	50	38	
		10	19	
		2	0	
		0.4	0	
Penconazole	EC 100 g/l	50	67	
		10	37	
		2	35	
		0.4	6	
Prochloraz	EC 400 g/l	50	18	
		10	1	
		2	0	
		0.4	0	
Pyrifenox	WP 500 g/kg	50	20	
		10	1	
		2	0	
		0.4	0	
Triadimefon	WP 250 g/kg	50	47	
		10	23	
		2	6	
		0.4	0	
Triadimenol	EC 250 g/l	50	68	
		10	45	
		2	21	
		0.4	1	
Spiroxamine	EC 500 g/l	10	2	
		2	0	
		0.4	0	
BP-4	Tankmix	10 + 10	84	72
Cyproconazole		0.4 + 0.4	23	19
BP-4	Tankmix	50 + 50	96	98
Dinocap		10 + 10	30	13
		2 + 2	2	0
		0.4 + 0.4	2	0
BP-4	Tankmix	50 + 50	100	99
Fenarimol		10 + 10	92	49
		2 + 2	49	34
		0.4 + 0.4	19	5

5	BP-4 Fenpropimorph	Tankmix	10 + 10	43	19
			2 + 2	6	0
			0.4 + 0.4	2	0
10	BP-4 Flusilazole	Tankmix	50 + 50	100	98
			10 + 10	57	29
			2 + 2	14	0
			0.4 + 0.4	1	0
15	BP-4 Penconazole	Tankmix	50 + 50	100	99
			10 + 10	96	45
			2 + 2	66	35
			0.4 + 0.4	23	6
20	BP-4 Prochloraz	Tankmix	50 + 50	100	98
			10 + 10	71	13
			2 + 2	11	0
			0.4 + 0.4	7	0
25	BP-4 Pyrifenox	Tankmix	50 + 50	100	98
			10 + 10	56	13
			2 + 2	7	0
			0.4 + 0.4	4	0
30	BP-4 Triadimefon	Tankmix	50 + 50	99	99
			10 + 10	39	32
			2 + 2	15	6
			0.4 + 0.4	2	0
35	BP-4 Triadimenol	Tankmix	50 + 50	100	99
			10 + 10	78	52
			2 + 2	30	21
			0.4 + 0.4	9	1
40	BP-4 Spiroxamine	Tankmix	10 + 10	37	15
			2 + 2	10	0
			0.4 + 0.4	2	0

50 **Example 15**

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Puccinia recondita* on wheat

55 **[0084]** The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XV:

Table XV

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	50	12	
		10	6	
		2	0	
		0.4	0	
Cyproconazole	SL 100 g/l	2	99	
		0.4	88	
Dinocap	WP 190 g/kg	50	54	
		10	23	
		2	13	
		0.4	4	
Fenarimol	SC 125 g/l	50	90	
		10	23	
		2	9	
		0.4	0	
Fenpropimorph	EC 750 g/l	50	60	
		2	15	
		0.4	2	
Flusilazole	WP 200 g/kg	50	100	
		2	28	
		0.4	5	
Penconazole	EC 100 g/l	50	26	
		10	10	
		2	3	
		0.4	0	
Prochloraz	EC 400 g/l	50	14	
		10	1	
		2	0	
		0.4	0	
Pyriphenox	WP 500 g/kg	50	6	
		10	1	

		2	0	
		0.4	0	
5	Triadimenol	EC 250 g/l	50	98
			10	69
			2	21
10			0.4	1
	Spiroxamine	EC 500 g/l	50	17
			10	4
			2	0
15	BP-4	Tankmix	2 + 2	100 99
	Cyproconazole		0.4 + 0.4	92 88
	BP-4	Tankmix	50 + 50	61 60
20	Dinocap		10 + 10	38 28
			2 + 2	26 13
			0.4 + 0.4	10 4
	BP-4	Tankmix	50 + 50	99 91
25	Fenarimol		10 + 10	52 28
			2 + 2	15 9
			0.4 + 0.4	12 0
	BP-4	Tankmix	50 + 50	84 65
30	Fenpropimorph		2 + 2	19 15
			0.4 + 0.4	13 2
	BP-4	Tankmix	50 + 50	100 100
35	Flusilazole		2 + 2	55 28
			0.4 + 0.4	14 5
	BP-4	Tankmix	50 + 50	78 35
40	Penconazole		10 + 10	19 16
			2 + 2	7 3
			0.4 + 0.4	4 0
	BP-4	Tankmix	50 + 50	34 25
45	Prochloraz		10 + 10	12 7
			2 + 2	4 0
			0.4 + 0.4	0 0
	BP-4	Tankmix	50 + 50	31 18
50	Pyrifenoxy		10 + 10	15 7
			2 + 2	5 0
			0.4 + 0.4	4 0

BP-4	Tankmix	50 + 50	98	98
Triadimenol		10 + 10	89	71
		2 + 2	39	21
		0.4 + 0.4	10	1
BP-4	Tankmix	50 + 50	40	27
Spiroxamine		10 + 10	11	10
		2 + 2	3	0

Example 16

Fungicidal efficacy of the mixture of BP-4 + other fungicides (4 day residual) against *Pyrenophora teres* on barley

[0085] The tank mixes were obtained from technical material of BP-4 and different formulations of different active ingredients. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XVI.

Table XVI

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	50	11	
		10	0	
		2	0	
		0.4	0	
Cyproconazole	SL 100 g/l	10	1	
		2	4	
Dinocap	WP 190 g/kg	50	22	
		10	5	
		2	0	
		0.4	0	
Fenarimol	SC 125 g/l	10	4	
		2	0	
Fenpropimorph	EC 750 g/l	10	3	
		2	0	
		0.4	0	
Flusilazole	WP 200 g/kg	50	48	

		10	26	
		2	8	
5	Prochloraz	EC 400 g/l	10	48
		2	19	
		0.4	8	
10	Pyrifenox	WP 500 g/kg	50	15
		10	8	
		10	1	
		2	0	
		0.4	0	
15	Spiroxamine	EC 500 g/l	50	16
		10	5	
		2	3	
20		0.4	0	
	BP-4	Tankmix	10 + 10	9 1
	Cyproconazole		2 + 2	10 4
25	BP-4	Tankmix	50 + 50	29 31
	Dinocap		10 + 10	15 5
			2 + 2	5 0
			0.4 + 0.4	9 0
30	BP-4	Tankmix	10 + 10	6 4
	Fenarimol		2 + 2	1 0
	BP-4	Tankmix	10 + 10	4 3
	Fenpropimorph		2 + 2	5 0
35			0.4 + 0.4	5 0
	BP-4	Tankmix	50 + 50	70 54
	Flusilazole		10 + 10	49 26
			2 + 2	26 8
40	BP-4	Tankmix	10 + 10	49 48
	Prochloraz		2 + 2	30 19
			0.4 + 0.4	19 8
45	BP-4	Tankmix	50 + 50	26 25
	Pyrifenox		10 + 10	10 8
	BP-4	Tankmix	50 + 50	29 20
	Triadimefon		10 + 10	10 5
50	BP-4	Tankmix	50 + 50	46 21
	Triadimenol		10 + 10	19 1

		2 + 2	3	0
		0.4 + 0.4	1	0
BP-4	Tankmix	50 + 50	43	26
Spiroxamine		10 + 10	24	5
		2 + 2	6	3
		0.4 + 0.4	3	0

Example 17

Fungicidal efficacy of the mixture of different benzophenones + metconazole (2 day curative) against *Blumeria graminis* f. sp. *tritici* on wheat

[0086] The tank mixes were obtained from technical material of the benzophenones BP-2 and BP-4 and metconazole. The benzophenones, the type of formulations, the observed and expected efficacies with different rates are given in Table XVII:

Table XVII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-2	EC 100 g/l	54	67	
		18	25	
		6	18	
		2	13	
		0.67	4	
BP-4	TC 100%	54	92	
		18	78	
		6	42	
		2	27	
		0.67	9	
Metconazole	EC 100 g/L	27	40	
		9	14	
		3	7	
		1	2	
		0.33	0	
BP-2 Metconazole	Tankmix	27 + 54	90	80
		9 + 18	67	35
		3 + 6	46	23
		1 + 2	22	12
		0.33 + 0.67	9	4

EP 1 023 834 A1

Table XVII (continued)

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4 Metconazole	Tankmix	27 + 54	98	95
		9 + 18	93	81
		3 + 6	49	45

Example 18

Fungicidal efficacy of the mixture of BP-4 + AP-4 (1 day residual) against *Uncinula necator* on vines

[0087] The tank mixes were obtained from technical material of BP-4 and AP-4. The active ingredients, the type of formulations, the observed and expected efficacies with different rates are given in Table XVIII: Table XVIII

Compound	Formulation	dose rate (ppm)	Observed Efficacy	Expected Efficacy
BP-4	TC 100 %	8	42	
		4	43	
		2	9	
		1	28	
		0.5	21	
		0.25	0	
AP-4	TC 100 %	1	76	
		0.5	76	
		0.25	50	
		0.12	32	
		0.06	7	
		0.03	13	
BP-4 AP-4	Tank mix	8 + 1	88	86
		4 + 0.5	91	87
		2 + 0.25	85	79
		1 + 0.12	89	83
		0.5 + 0.06	88	81
		0.25 + 0.03	79	76

B Field Tests

Example 19

Fungicidal efficacy of the mixture of BP-1 + metconazole in the field against the sugar beet disease *Cercospora beticola*

[0088] The tank mix was obtained from a SC formulation containing 100 g of BP-1 per liter and a SL formulation containing 60 g of metconazole per liter. The observed and expected efficacies are given in Table XIX


Table XIX

dose rate g / ha		Observed Efficacy	Expected Efficacy
BP-1	metconazole		
250	0	42.5	--
0	90	58	--
250	90	85.4	75.9

Claims

1. A fungicidal composition comprising a fungicidally acceptable carrier and/or surface active agent together with, and synergistically effective amounts of

(a) at least one benzophenone of formula I




wherein

- R¹ represents a halogen atom, an optionally substituted alkyl, alkanoyloxy or alkoxy group; or a hydroxy group,
 R² represents a halogen atom or an optionally substituted alkyl group,
 m is 0 or an integer of 1 to 3;
 R³ independently represents a halogen atom, an optionally substituted alkyl or alkoxy group or a nitro group;
 R⁴ represents a halogen atom, a cyano, carboxy, hydroxy or nitro group or an optionally substituted alkyl, alkoxy, alkenyl, alkylthio, alkylsulphinyl, alkylsulphonyl or amino group;
 R⁵ represents an optionally substituted alkyl group;
 R⁶ represents a halogen atom or a nitro group, an optionally substituted alkyl, alkoxy, alkenyloxy, alkynyloxy, alkylthio, cycloalkyl, cycloalkyloxy, aryloxy group;
 n is 0 or 1; and
 R⁷ independently represents a halogen atom, an optionally substituted alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy, cycloalkyl, cycloalkoxy group;

(b) at least one fungicidally active ingredient selected from the following groups (A), (B), (C), (D) and (E):


- (A) an ergosterol biosynthesis inhibitor;
 (B) a strobilurine derivative,
 (C) a melanin biosynthesis inhibitor,
 (D) a compound selected from the group consisting of acibenzolar, benomyl, captan, carboxin, chlorothalonil, copper, cyprodinil, dinocap, dithianon, dimethomorph, dodine, ethirimol, famoxadone, fenpiclonil, fluazinam, mancozeb, metataxyl, pyrifenoxy, sulfur, vinclozolin and
 (E) an azolopyrimidine of formula II



in which

- R^8 and R^9 each independently represent hydrogen or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, aryl, heteroaryl, cycloalkyl, bicycloalkyl or heterocyclyl group, or together with the interjacent nitrogen atom represent an optionally substituted heterocyclic ring,
- R^{10} represents hydrogen or an alkyl or aryl group,
- R^{11} represents a hydrogen or halogen atom or an alkyl or alkoxy group,
- L independently represents a halogen atom or an optionally substituted alkyl or alkoxy group,
- A represents N or CR^{12} , wherein R^{12} has the meaning given for R^{10} , and
- p is 0 or an integer from 1 to 5.

2. A composition as claimed in claim 1, wherein the benzophenone is a compound of formula IA,



wherein

- R^1 represents a halogen atom, a methyl, trifluoromethyl, methoxy or hydroxy group;
- R^2 represents a halogen atom or a methyl group;
- R^3 represents a bromo or chloro atom, a methyl, trifluoromethyl or nitro group;
- R^4 represents a methyl group;
- R^5 represents an alkyl group;
- R^6 and R^7 each independently represent an alkoxy group which may be substituted by a phenyl, alkylphenyl or halophenyl group; and
- n is 0 or 1.


3. A composition as claimed in claim 1, wherein the benzophenone of formula I is selected from

- 6'-butoxy-2,6-dichloro-4',5'-dimethoxy-2'-methylbenzophenone;
- 2,6-dichloro-4',5'-dimethoxy-6'-(2-fluorobenzyloxy)-2'-methylbenzophenone;
- 6'-benzyloxy-4',5'-dimethoxy-2,6-dimethyl-2'-methylbenzophenone;
- 3-bromo-2',6-dimethyl-2,4',5',6'-tetramethoxybenzophenone; and
- 2,6-dichloro-2'-methyl-4',5',6'-trimethoxybenzophenone.

4. A composition as claimed in claim 1, wherein the ergosterol biosynthesis inhibitor of group (A) is selected from fenarimol, fenpropimorph, fenpropidine, spiroxamine, triforine, cyproconazole, epoxiconazole, flusilazole, metco-

nazote, myclobutanil, penconazole, prochloraz, propiconazole, tebuconazole, triadimefon and tridimenol,

5. A composition as claimed in claim 4, wherein the ergosterol biosynthesis inhibitor of group (A) is metconazole.
6. A composition as claimed in claim 1, wherein the strobilurine derivative of group (B) is selected from azoxystrobin, kresoxim-methyl and trifloxystrobin.
7. A composition as claimed in claim 1, which comprises one compound of formula I, one strobilurine compound selected from group (B) and one ergosterol biosynthesis inhibitor of group (A), or one compound of formula I and two different compounds of group (D).
8. A composition as claimed in claim 1, wherein the melanin biosynthesis inhibitor of group (C) is selected from capropamid, chlobenthiazole, diclocymet, pyroquilon, phthalide, tricyclazote and AC 382042.
9. A composition as claimed in claim 1, wherein the azolopyrimidine of group (E) is a compound of formula IIA,



wherein

R^8 represents an alkyl, alkenyl, cycloalkyl or haloalkyl group, and
 R^9 represents a hydrogen atom, or an alkyl group; or
 R^8 and R^9 together form an optionally substituted alkylene group having 3 to 6 carbon atoms in the main chain, in which one CH_2 group may be replaced by O, S or NH;
 L^1 , L^2 and L^3 each independently represent a hydrogen or halogen atom or a C_{1-4} alkoxy group, at least one of which represents a halogen atom; and
Hal denotes a halogen atom.

10. A composition as claimed in claim 9, wherein Hal represents a chlorine atom, L^1 represents a fluorine atom; L^2 represents a hydrogen or fluorine atom or a methoxy group and L^3 represents a fluorine or chlorine atom.
11. A composition as claimed in claim 9, wherein the azolopyrimidine of formula II is selected from the group consisting of

5-chloro-6-(2-chloro-6-fluorophenyl)-7-(cyclopropylamino)-[1,2,4]triazolo[1,5-a]pyrimidine;
 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(4-methylpiperid-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine;
 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(2,2,2-trifluoroethylamino)-[1,2,4]triazolo[1,5-a]pyrimidine; and
 5-chloro-6-(2,4,6-trifluorophenyl)-7-[2-(1,1,1-trifluoro)propylamino]-[1,2,4]triazolo[1,5-a]pyrimidine.

12. A composition as claimed in claim 1, wherein the ratio (by weight) of the benzophenone of formula I to the fungicidal compound selected from the groups (A), (B), (C), (D) and (E) is from 10 : 1 to 1 : 10.
13. A composition as claimed in claim 12, wherein the ratio (by weight) of the benzophenone of formula I to the fungicidal selected from the groups (A), (B), (C), (D) and (E) is from 5 : 1 to 1 : 5.
14. Method of controlling the growth of fungi at a locus which comprises applying a composition as claimed in claim 1 to the locus.
15. Method of controlling the growth of powdery mildew at a locus which comprises applying a composition as claimed in claim 1 to the locus.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 30 0637

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
P,X	EP 0 933 025 A (AMERICAN CYANAMID CO) 4 August 1999 (1999-08-04) * page 2, line 54 - page 3, line 44 * * page 3, line 49 - page 4, line 29 * * page 7, line 8 - line 13 * * claims 6,11 *	1-15	A01N35/04 //(A01N35/04, 59:20,59:02, 55:00,47:44, 47:38,47:14, 47:04,43:90, 43:84,43:82, 43:76,43:653, 43:60,43:56, 43:54,43:42, 43:40,43:36, 43:32,43:30, 37:50,37:46, 37:38,37:34, 37:06)
A	EP 0 727 141 A (AMERICAN CYANAMID CO) 21 August 1996 (1996-08-21) * page 2, line 24 - page 3, line 1 * * page 11, line 51 - page 12, line 1 *	1-15	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A01N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 10 May 2000	Examiner Lamers, W
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 0637

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0933025 A	04-08-1999	JP 11269006 A	05-10-1999
EP 0727141 A	21-08-1996	AU 4209196 A	01-08-1996
		BR 9600165 A	06-01-1998
		CA 2167550 A	21-07-1996
		CN 1134929 A	06-11-1996
		CZ 9600089 A	14-08-1996
		HU 9600116 A	28-11-1996
		JP 8277243 A	22-10-1996
		SK 7496 A	06-11-1996
		US 5679866 A	21-10-1997
		US 5866722 A	02-02-1999